



IMPROVING SOIL HEALTH THROUGH ORGANIC AGRICULTURE: EFFECTS ON PHYSICOCHEMICAL AND BIOLOGICAL PROPERTIES

Mahnoor Ansari^{*1}, Fatima Roomesha², Hassam Sohail³

^{*1,2,3}Department of Agronomy, Faculty of Agriculture, University of Karachi, Karachi, Pakistan

^{*1}Mahnooransari85@gmail.com

Keywords

Organic agriculture;
Physicochemical properties;
Biological properties; Soil health;
Sustainable farming; Wheat

Article History

Received: 16 April 2025

Accepted: 22 May 2025

Published: 30 June 2025

Copyright @Author

Corresponding Author: *

Mahnoor Ansari

Abstract

Soil health is fundamental to sustainable agriculture and global food security. Organic farming practices are increasingly recognized for their ability to improve soil quality and crop productivity. This study assessed the impact of organic amendments—farmyard manure (FYM) and vermicompost (VC)—compared to conventional inputs, urea (U) and diammonium phosphate (DAP), on soil health indicators and wheat growth. The results showed significant ($P < 0.05$) enhancements in both soil physicochemical and biological properties under organic treatments relative to conventional fertilization. Organic amendments markedly increased soil organic matter (6.33-fold with VC and 5.70-fold with FYM), porosity (2.60-fold and 1.78-fold), and moisture content (1.93-fold and 1.81-fold). Conversely, bulk density and pH decreased by 1.19-fold and 1.12-fold under VC and FYM, respectively. Biological indicators, including microbial biomass carbon (MBC), nitrogen (MBN), and phosphorus (MBP), were also significantly elevated, showing 2.18-, 3.64-, and 4.10-fold increases with VC and 2.01-, 2.93-, and 3.08-fold with FYM. Additionally, populations of bacteria, fungi, and actinomycetes increased substantially under organic amendments. Correlation analyses highlighted strong interconnections between physicochemical and biological properties, emphasizing their collective role in sustaining soil health. Overall, organic amendments significantly enhanced soil fertility, microbial diversity, and biological activity compared to conventional fertilizers, reinforcing the critical role of organic agriculture in maintaining long-term soil productivity and ecological balance.

INTRODUCTION

Soil is a vital natural resource that supports agriculture, ecological balance, and environmental conservation (Hassan *et al.* 2022a). The essential properties of soils include their ability to support plant development, water, nutrients, and species, thus is central to sustainable agricultural system (Hore and Shoma 2024). But the latest techniques of integrated agricultural production involving intensive use of chemical inputs has made the situation worse with regard to soil, organic matter, and biological status (Hassan *et al.* 2022b). This

disturbing practice requires the examination of other forms of agriculture; like organic agriculture to remedy and rebuild suitable soil quality (Krif *et al.* 2020). Organic agriculture founded on the principles of using organic material, biological activities and farm ecosystem to support production (Hassan and David 2018). Organic farming uses organic substance such as farm yard manure, vermiculture compost, green manure and crop residues (Kwiatkowski *et al.* 2020). These inputs do not only deliver essential nutrient to the crops, but



also enhance the physico-chemical and biological property of the soil (Hassan *et al.* 2017). Modern technology has developed organic farming as a method of combating the adverse impacts of the traditional type of farming (Aulakh *et al.* 2022). These adverse impacts can be alleviated by enhancing soil biological resources, the proportion of organic materials in the soil, and by reducing pollution (Aulakh *et al.* 2022).

Most studies showed that soil health and its fertility depending on physicochemical indicators including bulk density (BD), porosity, organic matter (OM), moisture content and pH (Basak *et al.* 2020). It has been observed that FYM and VC has significantly enhanced the physical structure of the soil, water holding capacity (moisture), porosity and nutrients retention (Sharma, 2022). It has been examined that addition of FYM and VC significantly decreases the bulk density (BD) and pH, increases porosity, improves moisture and overall soil health index (Sivagamy *et al.* 2024). Organic agricultural practices use FYM and VC which served as source of food and energy for the microorganisms that are a major contributor in improving the OM, nutrient cycling and overall soil ecology (Sahu, 2023). Further, indices like microbial biomass carbon (MBC), nitrogen (MBN) and phosphorus (MBP), and community variability i.e. bacteria, fungi and actinomycetes are important attributes of soil quality (Basak *et al.* 2020). Organic agricultural practices i.e. use of FYM and VC enhance microbial biomass and numbers through supply of organic substances (Gao *et al.* 2022). This improves the soil biological properties and overall soil ecology and health (Gao *et al.* 2022). Plant growth promoting microorganisms include bacteria, fungi, and actinomycetes are involved in nutrient mineralization, organic matter decomposition and biological control against diseases (Shah *et al.* 2021). The application of organic amendments such as VC and FYM not only increases the number of soil microorganisms, but also increases microbial biomass, and physico-chemical indexes (Sharma, 2022).

In contrast the traditional agricultural practices i.e. use of urea (U) and diammonium phosphate (DAP) has been linked to negative effects on soil physico-chemical and biological properties in the long-run (Fellet *et al.* 2022). Physico-chemical and biological

properties, along with long-term soil health, are significantly enhanced by organic practices (Fellet *et al.*, 2022). In contrast, traditional practices, i.e. use of synthetic fertilizers (urea and DAP), often have transient positive effects and adverse long-term consequences (Kumar *et al.*, 2024). The effect of traditional agricultural practices (Urea and DAP) on the soil microbial biomass and community is negative, which kept aggravating with the time (Ramazanoglu *et al.* 2024). Traditional practices (Urea and DAP) always have detrimental impacts on the soil health and overall agro ecology (Teka *et al.* 2024). Arjjumend *et al.* (2021) concluded that effect of synthetic fertilizers is negative on the long-run. Omer *et al.* (2024) found that traditional practices i.e. use of synthetic chemicals and tillage have detrimental effects on the soil physical conditions and biological diversity. Despite growing evidence of the benefits of organic farming, further research is needed to quantify its effects on soil health indicators i.e. physical, chemical and biological properties. This study aims to fill this gap by analyzing firstly to assess the impact of organic practices on soil physico-chemical properties. Secondly to analyze its potential role in enhancing biological properties and microbial community and diversity of soil.

1. Material and Methods

1.1. Soil sampling

Samples (0-30 cm depth) were collected by using a hand auger. Field-moist soil samples were sieved (<2 mm) and divided into two subsamples. One subsample was air-dried at room temperature and used for chemical analyses. Second subsample was stored at 4°C for the analysis of biological properties.

1.2. Experimental setup

Five treatments (Table 1) were performed in triplicate, following a randomized complete block design. Fifteen pots (8 × 12 cm) were filled with 5 kg of air-dried and sieved (2 mm) soil. Treatments i.e. VC (T1), FYM (T2), U (T3) and DAP (T4) were applied at the rate of 30 g kg⁻¹ soil (Lhamu *et al.* 2024 and Ilker *et al.* 2016). After 60 days samples were collected from each pot for the analysis of physico-chemical properties i.e. OM, porosity, BD, moisture, pH and



EC and biological properties i.e. MBC, MBN, and MBP and bacteria, fungi and actinomycetes respectively.

1.3. Physico-chemical properties of soil

Organic matter was determined by measuring convertible organic carbon contents (Hassan *et al.* 2014). Porosity was measured through soil bulk and particle density (Rehman *et al.* 2008). Moisture contents were assessed via gravimetric method by weighing the soil before and after oven-drying (MBN), and phosphorus (MBP) via pre-fumigated and un-

(Hassan *et al.* 2014). The pH and EC were examined by using a soil-water ratio of 1:5 with a calibrated and properly rinsed pH and EC meter (Hassan *et al.* 2014). Basic physico-chemical properties of experimental soil are given in Table 2.

1.4. Biological properties of soil

The method of fumigation-extraction was used to determine the microbial biomass carbon (MBC), nitrogen

Table 1. Treatments of the experiment

| Treatments | Level (g kg ⁻¹ soil) | Rate (g kg ⁻¹) |
|------------|---------------------------------|----------------------------|
| T1 | VC | 30 |
| T2 | FYM | 30 |
| T3 | U | 30 |
| T4 | DAP | 30 |
| T5 | Control (CK) | 0 |

fumigated soil samples (Hassan *et al.* 2014). Whereas, microbial population i.e. bacteria, fungi and actinomycetes were quantified through serial dilution and culturing on selective media plates, and incubating under pretermitted conditions (Hassan and David, 2018).

2. Statistical analysis

The ANOVA test was used to analyze the collected data, while LSD test was employed to estimate the differences among treatments.




Figure 3. Pearson correlation heat map between the parameters



3. Results

3.1. Effect of organic versus traditional agricultural practices on the physico-chemical properties of soil

The effect of organic versus traditional agricultural practices on the physico-chemical properties of soil is given in Table 1. The organic inputs i.e. VC and FYM significantly ($P < 0.05$) increased all the measured soil physico-chemical properties compared to traditional practices and control. The increase in the OM via VC and FYM was 6.33-fold and 5.70-

fold higher compared to control. Conversely, the increase in the OM via traditional practices, i.e. DAP and urea was 1.78-fold and 1.60-fold higher compared to control. Similarly, increase in porosity through organic practices i.e. VC and FYM was 2.60-fold and 1.78-fold compared to control. Whereas the increase in the traditional practices i.e. DAP and urea was 1.28-fold and 1.16-fold compared to control. On the contrary, in organic practices i.e. VC and FYM a decrease in the BD was observed i.e. 1.19-fold and 1.14-fold. Whereas

fold, but this increase was significantly lower compared to organic practices. On the other hand, traditional practices increased the soil pH i.e. 1.12-fold and 1.08-fold compared to control. Whereas organic practices decreased the soil pH i.e. 1.19-fold and 1.12-fold compared to control.

fold, but this increase was significantly lower compared to organic practices. On the other hand, traditional practices increased the soil pH i.e. 1.12-fold and 1.08-fold compared to control. Whereas organic practices decreased the soil pH i.e. 1.19-fold and 1.12-fold compared to control.

3.2. Effect of organic versus traditional agricultural practices on the biological properties of soil




Figure 1. Effect of organic and traditional agricultural practices on the microbial biomass MBC, microbial biomass carbon, MBN, microbial biomass nitrogen, MBP, microbial biomass phosphorus, VC, vermicompost, FYM, farmyard manure; DAP, diammonium phosphate.

The effect of organic versus traditional agricultural practices on the biological properties of soil is given in Table 2. The organic inputs i.e. VC and FYM significantly ($P < 0.05$) increased all the measured biological soil properties compared to traditional practices and control. The increase in the MBC, MBN and MBP via VC i.e. 2.18-fold, 3.64-fold and 4.1-fold FYM i.e. 2.01-fold, 2.93-fold and 3.08-fold higher compared

to control. Conversely, the increase in the MBC, MBN and MBP via traditional practices i.e. DAP and urea was 1.48-fold, 1.76-fold and 2.03-fold and 1.24-fold, 1.29-fold and 1.42-fold higher compared to control. Similarly, increase in microbial population i.e. bacteria, fungi and actinomycetes through organic practices i.e. VC and FYM was 3.45-fold, 6.36-fold and 4.0-fold and 3.45-fold, 6.36-



fold and 4.0-fold respectively compared to control. Whereas the increase in the traditional practices, i.e., DAP and urea, was 1.91-fold, 1.36-fold and 2.0-fold and 1.36-

fold, 1.81-fold and 1.33-fold correspondingly compared to control.

3.3. Correlation analysis




Figure 2. Effect of organic and traditional agricultural practices on the microbial density CFU, colony forming unit; VC, vermicompost; FYM, farm yard manure; DAP,

Table 2. Basic physico-chemical properties of experimental soil

| Properties | Reading |
|------------------------------------|------------|
| Clay (g kg ⁻¹) | 100.6 |
| Silt (g kg ⁻¹) | 176.3 |
| Sand (g kg ⁻¹) | 723.1 |
| Textural class | Sandy loam |
| pH (1:2.5) | 7.98 |
| EC (µS cm ⁻¹) | 11.5 |
| CEC (C molc Kg ⁻¹) | 6.49 |
| OM (%) | 0.51 |
| Available N (mg kg ⁻¹) | 5.21 |
| Available P (mg kg ⁻¹) | 3.61 |
| Available K (mg kg ⁻¹) | 97.4 |
| Total P (g kg ⁻¹) | 0.21 |
| Total N (g kg ⁻¹) | 0.32 |

biological properties of the soil is given in Fig. 3. It shows that both physico-chemical and biological indices are closely interlinked with each other. been

observed that among physico-chemical Correlation between the physico and properties OM (0.847), pH (0.705) and EC (0.787) showed highly significant correlation with other



physiochemical indices and also with biological indices.

Whereas, among biological indices MBC (0.833), MBN (0.751), MBP (0.847) and bacteria (0.681) and fungi (0.658) showed a highly significant correlation with other biological and physiochemical indices.

4. Discussion

The effect of traditional i.e. U and DAP and organic i.e. VC and FYM agricultural practices on soil physico-chemical properties i.e. OM, porosity, BD, moisture, pH and EC are given in. The use of organic agricultural practices i.e. VC and FYM significantly ($P < 0.05$) improved all the measured properties compared to traditional agricultural practices and control. The maximum increase in OM, porosity, moisture, pH and EC and maximum decrease in BD was observed at VC and then in FYM. Conversely the effect of traditional or non-agricultural practices i.e. U and DAP was significantly ($P < 0.05$) low compared to agricultural practices and control. Overall effect of treatments was in order VC < FYM < DAP

< U < CK. Basak *et al.* (2020) found that organic inputs significantly improved soil physicochemical properties, including moisture, porosity, OM, and EC, enhancing overall soil health. Kindle *et al.* (2024) emphasized OM, porosity, BD, moisture, pH, and EC as key soil health indicators, highlighting the potential of organic practices to improve them. Sharma (2022) reported that VC and FYM notably enhanced OM, porosity, moisture, nutrient availability, and retention compared to traditional practices. Sivagamy *et al.* (2024) reported that VC and FYM significantly increased porosity, moisture, EC, and root

penetration, while reducing BD, pH, and overall soil health index. Sahu (2023) found that properties i.e. MBC, MBN and MBP and bacteria, fungi and actinomycetes are given in table Fig. 1-2. The use of organic agricultural practices i.e. VC and FYM significantly ($P < 0.05$) improved all the measured biological properties compared to traditional agricultural practices and control. The maximum increase in MBC, MBN and MBP and bacteria, fungi and actinomycetes was observed at VC and then in FYM. Conversely the effect of traditional or non-agricultural practices i.e. U and DAP was significantly ($P < 0.05$) low compared to agricultural practices and control. Overall effect of treatments was in order VC < FYM < DAP

< U < CK. Basak *et al.* (2020) highlighted microbial biomass (MBC, MBN, MBP) and community diversity (bacteria, fungi, and actinomycetes) as key soil quality indicators, with organic practices positively influencing them. Gao *et al.* (2022) found that FYM and VC significantly increased microbial biomass and improved soil biology and ecology. Shah *et al.* (2021) emphasized that plant growth-promoting microorganisms aid in nutrient mineralization, organic matter decomposition, and disease control. Sharma (2022) reported that VC and FYM enhance microbial populations, biomass, and soil fertility, boosting crop yield. In contrast, Fellet *et al.* (2022), Ramazanoglu *et al.* (2024), and Teka *et al.* (2024) found that synthetic fertilizers (Urea and DAP) negatively impact microbial biomass, community structure, and enzyme activity, with effects worsening over time, ultimately harming soil ecology and agro-ecosystems. It has been examined that synthetic fertilizers have momentary positive effects but in the long-run highly negative impacts on the

Table 3. Effect of organic and traditional agricultural practices on the physico-chemical properties of soil

| Treatment | OM (%) | Porosity (%) | BD (g/cm ³) | Moisture (%) | pH (1:1.5) | EC (1:1.5) |
|-----------|------------|--------------|-------------------------|--------------|-------------|------------|
| VC | 3.23±0.23a | 66±1.41a | 0.92±0.05e | 62±1.87a | 7.23±0.86d | 10.2±0.56a |
| FYM | 2.91±0.19b | 57±1.21b | 0.97±0.04d | 58±1.56b | 7.58±0.72c | 8.61±0.45b |
| DAP | 0.91±0.12c | 41±1.11c | 1.04±0.03bc | 43±1.11c | 7.87±0.52bc | 6.01±0.31c |
| Urea | 0.82±0.11d | 37±1.01d | 1.08±0.02ab | 38±1.01d | 7.94±0.41ab | 4.04±0.26d |



| | | | | | | |
|---------|------------|----------|------------|----------|------------|------------|
| Control | 0.51±0.09e | 32±0.91e | 1.10±0.01a | 32±0.93e | 7.98±0.32a | 3.34±0.21e |
|---------|------------|----------|------------|----------|------------|------------|

organic practices improve soil properties and health by providing nutrients and energy for microorganisms, enhancing OM, carbon, and nutrient cycling. In contrast, Fellet *et al.* (2022) and Kumar *et al.* (2024) observed that synthetic fertilizers (Urea and DAP) negatively impact soil physico-chemical properties, with mostly transitory or harmful long-term effects.

The effect of traditional i.e. U and DAP and organic i.e. VC and FYM agricultural practices on soil biological overall soil health (Arjjumend *et al.* 202; Omer *et al.* 2024).

It has been observed that among the physico-chemical properties, organic matter (OM) (0.847), pH (0.705), and electrical conductivity (EC) (0.787) exhibit highly significant correlations with other physico-chemical indices as well as biological indices. Basak *et al.* (2020), in a study, demonstrated that soil health and fertility depend on physico-chemical indicators such as porosity, OM, EC, and pH, which are closely interrelated. Among the biological indices, microbial biomass carbon (MBC) (0.833), microbial biomass nitrogen (MBN) (0.751), microbial biomass phosphorus (MBP) (0.847), bacteria (0.681), and fungi (0.658) showed highly significant correlations with other biological and physico-chemical indices. Indices such as MBC, MBN, and MBP, along with community variability (e.g., bacteria, fungi, and actinomycetes), are critical attributes of soil quality and are interrelated with each other as well as with other indices (Basak *et al.*, 2020). Similarly, in an experiment, Shah *et al.* (2021) identified a significant ($P < 0.05$) correlation among physico-chemical and biological indices and further concluded that all these indices play a vital role in maintaining the soil health index.

5. Conclusion

In conclusion, organic agriculture practices demonstrate significant potential to enhance soil physico-chemical and biological properties. Thereby improving long-term soil health and food security, unlike traditional inorganic practices that offer only temporary benefits. Given the strong interconnections between soil properties (e.g., OM,

pH and EC) and biological indices (e.g., MBC, MBN, MBP, bacteria and fungi), transitioning to organic farming is essential for maintaining a healthy and sustainable soil ecosystem, offering valuable insights for farmers, researchers, and policymakers.

Funding

Funded by Chinese Academy of Sciences President, s International Fellowship Initiative Grant No. 2023VCC0007. **Acknowledgment**

Author also acknowledges Chinese Academy of Sciences Presidents International Fellowship Initiative is acknowledged for funding.

Conflict of Interest

The authors declare no conflict of interest.

References

- Arjjumend, H., Koutouki, K., & Neufeld, S. (2021). Comparative advantage of using biofertilizers in indian agroecosystems: an analysis from the perspectives of stakeholders. *European Journal of Agriculture and Food Sciences*, 3: 26-36.
- Aulakh, C.S., Sharma, S., Thakur, M., & Kaur, P. (2022). A review of the influences of organic farming on soil quality, crop productivity and produce quality. *Journal of Plant Nutrition* 45: 1884-1905.
- Basak, B. B., Saha, A., Sarkar, B., Kumar, B. P., Gajbhiye, N. A., & Banerjee, A. (2021). Repurposing distillation waste biomass and low-value mineral resources through biochar-mineral-complex for sustainable production of high-value medicinal plants and soil quality improvement. *Science of The Total Environment* 760: 143319.
- Fellet, G., Conte, P., Bortolotti, V., Zama, F., Landi, G., Chillura Martino, D.F., Ferro, V., Marchiol, L., & Lo Meo, P. (2022). Changes in physicochemical properties of biochar after addition to soil. *Agriculture* 12: 320.



- Gao, D., Bai, E., Wang, S., Zong, S., Liu, Z., Fan, X., Zhao, C., & Hagedorn, F. (2022). Three-dimensional mapping of carbon, nitrogen, and phosphorus in soil microbial biomass and their stoichiometry at the global scale. *Global Change Biology* 28: 6728-6740.
- Hassan, W., Saba, T., Jabbi, F., Wang, B., Cai, A., & Wu, J. (2022a). Improved and sustainable agroecosystem, food security and environmental resilience through zero tillage with emphasis on soils of temperate and subtropical climate regions: A review. *International Soil and Water Conservation Research* 1: 530-545.
- Hassan, W., Saba, T., Wu, J., Bashir, S., Bashir, S., Gatashah, M.K., Diao, Z.H. & Chen, Z. (2022b). Temperature responsiveness of soil carbon fractions, microbes, extracellular enzymes and CO₂ emission: mitigating role of texture. *PeerJ* 10: 13151.
- Hassan, W., David, J. (2018) Comparative effectiveness of ACC-deaminase and/or nitrogen fixing rhizobacteria in rice (*Oryza sativa* L.). *Environmental Engineering and management Journal* 17: 1113-1121.
- Hassan, W., Bashir, S., Hanif, S., Sher, A., Sattar, A., Wasaya, A., & Hussain, M. (2017). Phosphorus solubilizing bacteria and growth and productivity of mung bean (*Vigna radiata*). *Pakistan Journal of Botany* 49: 331-336.
- Hore, S. (2024) Assessment of soil chemical characteristics in the context of Bangladesh: a comprehensive review. *Community and Ecology* 2: 1.
- Kindie, Z., Yedem Fentie, A., & Girma, M. (2023) Contribution of on-farm avocado (*Persea americana*) tree- based agroforestry practice on selected soil physical and chemical properties of Inguti small watershed, in the highlands of North-Western Ethiopia. *Sustainable Environment* 9: 2289702.
- Kumar, P., Basak, B.B., Patel, V.J., Senapati, N., Ramani, V.P., Gajbhiye, N.A. & Kalola, A.D. (2024) Enriched soil amendments influenced soil fertility, herbage yield and bioactive principle of medicinal plant (*Cassia angustifolia* Vahl.) grown in two different soils. *Heliyon* 10: 3.
- Krif, G., Mokri, F., Aissami, A.E., Laasli, S.E., Imren, M., Özer, G., Paulitz, T., Lahlali, R. & Dababat, A.A. (2020) Diversity and management strategies of plant parasitic nematodes in Moroccan organic farming and their relationship with soil physico-chemical properties. *Agriculture* 10: 447.
- Kwiatkowski, C., A., & Harasim, E. (2020) Chemical properties of soil in four-field crop rotations under organic and conventional farming systems. *Agronomy* 10: 1045.
- Lhamu, S., Paul, A., Boruah, A., Yedi, N., Alam, S., Kumar, M. and Zimik, Y.V., Impact of different organic nutrient sources on soil chemical properties in rabi-season rapeseed cultivation.
- Ilker, U.Z., Sonmez, S., Tavali, I.E., Citak, S., Uras, D.S., & Citak, S. (2016) Effect of vermicompost on chemical and biological properties of an alkaline soil with high lime content during celery (*Apium graveolens* L. var. *dulce* Mill.) production. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* 44: 280-290.
- Omer, E., Szlatenyi, D., Csenki, S., Alrwashdeh, J., Czako, I., & Láng, V. (2024) Farming practice variability and its implications for soil health in agriculture: a review. *Agriculture* 14: 2114.
- Rahman, M.H., Okubo, A., Sugiyama, S., & Mayland, H.F. (2008) Physical, chemical and microbiological properties of an Andisol as related to land use and tillage practice. *Soil and Tillage Research* 101: 10-19.
- Ramazanoglu, E., Beyyavas, V., Cevheri, C. İ., Sakin, E., & Yilmaz, S. N. (2024) Effects of farmyard manure and chemical fertilizer application rates on soil biology, cotton and fiber yield. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* 52: 13838-13838.
- Sahu, S.G. (2023) Soil properties under continuous application of inorganic and organic sources of nutrients in cereal-vegetable-pulse cropping system in an Inceptisols (Doctoral dissertation, Department of Soil Science and Agricultural Chemistry, OUAT, Bhubaneswar).



- Shah, K. K., Tripathi, S., Tiwari, I., Shrestha, J., Modi, B., Paudel, N., & Das, B. D. (2021) Role of soil microbes in sustainable crop production and soil health: A review. *Agricultural Science and Technology* 13: 109-118.
- Sharma, S.B. (2022) Trend setting impacts of organic matter on soil physico-chemical properties in traditional vis-a-vis chemical-based amendment practices. *PLOS Sustainability and Transformation* 1: 0000007.
- Singh, K. C., Chauhan, R. S., Goswami, G., Bhosale, T. A., & Haobijam, J. W. (2023) Evaluation of the possibility of establishing certain organic cultivation packages as nutrient sources for chamomile (*Matricaria chamomilla* L.) cultivation at temperate region of Uttarakhand, India. *International Journal of Environment and Climate Change* 13: 2167-2179.
- Sivagamy, K., Karunakaran, V., Ananthi, K., Rajesh, M., Prasad, S. A., Tamilselvi, C. & Kanaga, S. (2024) Effects of organic manures and foliar sprays on growth and yield of finger millet. *Agronomy Journal of Nepal* 8: 63-69.
- Teka, K., Abraha, B., Mebrahtom, S., Tsegay, A., Welday, Y., Gessesse, T. A., & Hansson, L. (2024) Effect of vermicompost on soil fertility and crop productivity in the dry lands of Ethiopia. *Compost Science & Utilization* 31: 75-8