



EFFICACY OF BIOPESTICIDES AGAINST SUCKING PESTS AND THEIR IMPACT ON BITTER GOURD (*MOMORDICA CHARANTIA* L.) YIELD

Kamran Ali¹, Tariq Mahmood Soomro^{*2}, Shahid Iqbal³

^{1, *2, 3}Institute of Agricultural Chemistry and Soil Science, Punjab Agricultural Research Council, Lahore, Pakistan

kamran.ali78@gmail.com, ²tariq.soomro814@gmail.com, ³shahid.iqbal523@gmail.com

Keywords

Bitter gourd, *Azadirachta indica*, Cypermethrin, Biopesticides, Sucking pests, Whiteflies, Thrips, Integrated Pest Management (IPM)

Article History

Received: 30 April 2026

Accepted: 15 June 2026

Published: 30 June 2026

Copyright @Author

Corresponding Author: *

Tariq Mahmood Soomro

Abstract

A field experiment was conducted at the Agricultural Research Institute, Tarnab, Peshawar, Pakistan, during 2017–2018 to evaluate the effectiveness of different botanical extracts in regulating the population dynamics of major sucking pests, including aphids, whiteflies, and thrips, in bitter gourd (*Momordica charantia* L.). The experiment was arranged in a Randomized Complete Block Design (RCBD) with three replications and consisted of five treatments: cypermethrin, *Azadirachta indica* (neem) extract, *Parthenium hysterophorus* extract, *Eucalyptus globulus* extract, and an untreated control.

The results demonstrated that both synthetic insecticide and botanical extracts significantly reduced pest populations and improved crop productivity. The lowest mean populations of aphids (0.88/leaf), whiteflies (0.90/leaf), and thrips (0.93/leaf) were recorded in cypermethrin-treated plots, followed by neem extract. In contrast, untreated control plots exhibited the highest pest infestations, with mean populations of 5.60 aphids/leaf, 8.89 whiteflies/leaf, and 5.96 thrips/leaf. Cypermethrin treatment produced the maximum fruit number (22.79 fruits per plant) and yield (24.80 kg per treatment), followed by neem extract, whereas the control treatment recorded the lowest yield (10.81 kg per treatment).

Among the tested botanical treatments, neem extract showed the highest effectiveness in suppressing sucking pest populations and enhancing yield compared with *Eucalyptus* and *Parthenium* extracts. These findings suggest that neem-based botanical formulations have considerable potential as an important component of Integrated Pest Management (IPM) programs for bitter gourd production, offering an environmentally sustainable approach for reducing reliance on synthetic insecticides.

Introduction

Bitter gourd (*Momordica charantia*), commonly known as karela, bitter melon, and balsam pear, is an essential vegetable crop in Southeast Asia, celebrated for its high nutritional and medicinal value. It belongs to the Cucurbitaceae family, alongside other important crops like squash, watermelon, muskmelon, and cucumber (Palada and Chang, 2003). Bitter gourd is rich in vitamins and minerals and is particularly known for its high

concentrations of Vitamin C and Vitamin A, which are significantly higher compared to other cucurbits (Palada and Chang, 2003). Additionally, it is widely recognized for its medicinal properties, particularly in managing diabetes, due to its bioactive components such as antioxidants (Krawinkel and Keding, 2006). Bitter gourd is not only a food source but also serves as a traditional medicine in many cultures for treating a variety of ailments, including digestive disorders, skin



diseases, and liver problems (Kubola and Siriamornpun, 2008). In terms of cultivation, bitter melon is an annual crop grown in tropical and subtropical regions. It is typically sown from January to June in regions with mild winters, although in colder climates, it is often grown as a summer crop (Singh et al., 2006). Bitter melon thrives in well-drained, sandy loam soil, and its cultivation requires moderate temperatures and sufficient sunlight. In Pakistan, bitter melon is cultivated on 6017 hectares, yielding approximately 56949 tons of produce. In the Khyber Pakhtunkhwa province, 528 hectares are dedicated to bitter melon cultivation, producing 4713 tons annually (MINFAL, 2018-19). The crop is an important part of the agricultural landscape in Pakistan, providing both food and income for many farmers. Despite its numerous benefits, it is affected by several pests, including aphids, whiteflies, thrips, fruit flies, and mites, which attack various stages of plant growth and cause significant yield losses. These pests are known to cause damage by feeding on the plant, leading to reduced photosynthesis, wilting, and in severe cases, the death of the plant (Hussain and Ahmad, 2012). The damage caused by these pests can lead to crop losses ranging from 30% to 80%, severely affecting the productivity of bitter melon farms. Aphids, such as *Myzus persicae*, are one of the most common and damaging pests in bitter melon cultivation. They are sap-sucking insects that feed on the leaves, young shoots, and stems, leading to wilting and deformation of plant parts. Aphids also excrete a sugary substance called honeydew, which encourages the growth of sooty mould on the leaves, further reducing the plant's ability to photosynthesize (Capinera, 2005; Barbercheck, 2014). Aphids are known to carry and transmit several plant viruses, including the Cucumber mosaic virus and the Tobacco mosaic virus, which can severely reduce plant vigour and yield. Thrips (*Thrips tabaci*) are another major pest affecting bitter melon. These tiny insects are known to damage flowers, leading to reduced fruit set and deformation of the plant's structure. Female thrips lay their eggs inside the flowers, and the larvae feed on the plant tissues, which can reduce flower production by up to 60% (GPP, 2009). Thrips can also transmit viral diseases, further

compounding the problem. The feeding activity of thrips leads to significant yield losses in bitter melon crops. Whiteflies (*Bemisia tabaci*) are also significant pests of bitter melon. These small, sap-sucking insects spend most of their life cycle on the underside of leaves. Whiteflies are known to cause damage to bitter melon by feeding on the plant's sap, leading to yellowing of the leaves, reduced plant vigour, and stunted growth. Whiteflies also excrete honeydew, which can cause the development of sooty mould on the leaves, further inhibiting photosynthesis (Mustafa, 1995; Nyoike, 2007). In addition to direct feeding damage, whiteflies are vectors for several plant viruses, including the Tomato yellow leaf curl virus, which can cause severe damage to crops (Abrahamian and Abou-Jawdah, 2014). Fruit flies, such as *Bactrocera cucurbitae*, are notorious for attacking the fruits of bitter melon. These pests lay their eggs inside the fruits, and the larvae feed on the fruit pulp, causing it to rot and leading to significant crop loss. The presence of fruit flies can make the harvested produce unmarketable, reducing the overall yield of bitter melon crops (Singh et al., 2006). To mitigate the damage caused by these pests, a variety of pest management strategies have been developed. Traditional pest management approaches include the use of chemical insecticides, cultural practices, and biological control methods. However, chemical insecticides, while effective, have significant drawbacks, including the development of resistance in pest populations and harmful effects on non-target organisms, including beneficial insects and humans (Sarwar, 2015). Moreover, many of these chemicals are not environmentally friendly and can accumulate in the food chain, posing risks to human health and the ecosystem (Kumar and Thakur, 2017). Cultural practices such as crop rotation, mulching, and the use of reflective mulches have been employed to reduce pest populations and improve crop health. Mulching, in particular, has been shown to reduce the infestation of whiteflies and other pests by preventing their colonization of the plant (Frank and Liburd, 2005). Other cultural



practices include the removal of infected plant debris and the use of sticky traps to pest populations (Basu, 1995). Biological control methods have also gained popularity as a more sustainable alternative to chemical pesticides. Natural enemies of aphids, such as ladybird beetles and parasitic wasps, can help suppress pest populations (Kos et al., 2008). Fungal pathogens like *Beauveria bassiana* and *Metarhizium anisopliae* have shown promise in controlling whiteflies and thrips, providing a more environmentally friendly pest management solution (Faria and Wraight, 2001).

Objective of the Study:

Given the challenges posed by pests in bitter gourd cultivation and the potential harm caused by chemical pesticides, this study aims to evaluate the effectiveness of natural plant extracts and a synthetic insecticide (Cypermethrin) in controlling the population density of aphids, whiteflies, and thrips in bitter gourd crops.

Results

The experiment regarding “effect of different plant extracts on the population density of aphids, whitefly and thrips on Bitter gourd *Momordica charantia* was conducted at the Agricultural Research Institute, Tarnab, Peshawar, Pakistan, during 2017-18. Randomized Complete Block Design (RCBD) was used in this trial, having five treatments (Cypermethrin ®25% EC (Sipermethrin 80-120

ml/ha), *Azadirachta indica* 10 Kg/ha, *Parthenium hysterophorus*. L. leaves extract 10 Kg/ha, *Eucalyptus globus* leaves extract 10 Kg/ha and Control (Water spray) that was replicated three times. Data on the relevant parameters are as follows. The experiment conducted in 2017-18 at the Agricultural Research Institute Tarnab, Peshawar, comparing Cypermethrin and various plant extracts (*Azadirachta indica*, *Parthenium hysterophorus*, *Eucalyptus globus*, and a control) on pest control in Bitter gourd, showed significant differences across treatments. Cypermethrin was the most effective, reducing aphid, whitefly, and thrips populations significantly, with the lowest mean values observed for pests and the highest fruit yield (22.79 fruits/plant and 20670 kg/ha). Among plant extracts, *Azadirachta indica* was the most effective botanical treatment, significantly reducing pest populations and resulting in 18.55 fruits/plant and 16126 kg/ha yield. *Parthenium hysterophorus*. And *Eucalyptus Globus* was less effective, with yields of 13411 kg/ha and 13374 kg/ha, respectively. The control treatment showed the highest pest populations (aphids: 5.60, whiteflies: 7.91, and thrips: 5.96), and the lowest yield (9008 kg/ha). These results highlight that while Cypermethrin provided the best control, *Azadirachta indica* performed comparably in pest management, making it a viable alternative in Integrated Pest Management (IPM) strategies.

Table: Comparison of Natural Plant Extracts and Cypermethrin on Different Parameters in Bitter Gourd Crop

<i>hyste ropho rus.</i>						
(T3)						
Eucal	3.22	2.82b	3.16	15.3	16.0	13
yptus	b		b	8c	4c	37
glob						4c
us						
(T4)						
Contr	5.60a	7.91a	5.96	9.62	10.8	90



ol (T5) a d 1d 08 d

The population density of aphids was significantly lower in the Cypermethrin-treated plots (0.88) compared to all the plant extracts and the control. The control had the highest aphid population (5.60). Statistically, the treatments showed significant differences, with Cypermethrin being the most effective in reducing aphid density. Cypermethrin also showed the lowest whitefly population (0.90), while the control had the highest (7.91). The results showed significant differences between the treatments, with Cypermethrin outperforming all plant extracts, followed by *Azadirachta indica* and *Parthenium hysterophorus*. Similarly, Cypermethrin was the most effective treatment in reducing thrips (0.93), while the control again showed the highest thrip population (5.96). Statistically significant differences were observed among the treatments. Cypermethrin-treated plants had the highest mean number of fruits per plant (22.79), followed by *Azadirachta indica* (18.55). The control plants had the lowest mean (9.62). Statistical differences were

observed among all treatments, with Cypermethrin showing the best result. Cypermethrin-treated plants had the highest yield per treatment (24.80 kg), significantly higher than the other treatments. The control plants had the lowest yield per treatment (10.81 kg). Statistical analysis confirmed that Cypermethrin was the most effective in terms of yield. The yield per hectare followed similar trends, with Cypermethrin having the highest yield (20670 kg/ha) and the control having the lowest (9008 kg/ha).

Treatment s	Aph ids/ Leaf	White flies/ Leaf	Thri ps/L eaf	Frui ts/Pl ant	Yiel d per Tre atment (kg)	Yi el d (k g/ ha)
Cype rmet hrin (T1)	0.88 d	0.90e	0.93 d	22.7 9a	24.8 0a	20 67 0a
<i>Azad irach ta indic a</i> (T2)	2.60c	2.18d	2.66 c	18.5 5b	19.3 5b	16 12 6b
<i>Parth eniu m</i>	3.00 b	2.66c	3.12 b	15.2 0c	16.0 5c	13 41 1c

Discussion

An experiment was conducted at the Agricultural Research Institute, Tarnab, Peshawar, Pakistan,

during 2017-18 to evaluate the effect of different plant extracts and Cypermethrin on the population density of aphids, whiteflies, and thrips on Bitter



gourd (*Momordica charantia*). The experiment utilized a Randomized Complete Block Design (RCBD) with five treatments: Cypermethrin (25% EC), *Azadirachta indica* (10 Kg/ha), *Parthenium hysterophorus*. (10 Kg/ha), *Eucalyptus globus* (10 Kg/ha), and Control (water spray), replicated three times. The results showed that Cypermethrin was the most effective in reducing the population of aphids, whiteflies, and thrips, leading to significantly higher fruit yield. This is because Cypermethrin attacks the insect nervous system and inhibits sodium ions, leading to rapid pest reduction (Rahman et al., 2014; Kumar and Thakur, 2017; Sharma and Tayde, 2017). These findings are consistent with Ali et al. (2011), who reported the effectiveness of chemical insecticides over natural extracts. Among the natural plant extracts, *Azadirachta indica* was the most effective against aphids, whiteflies, and thrips, resulting in higher fruit numbers and yield. Azadirachtin, the active compound in *Azadirachta indica*, has anti-feeding and growth-regulating properties, which hinder insect growth (Sarwar, 2015). This is supported by previous studies, including those of Kumar and Thakur (2017) and Kuswaha and Painka (2016). *Parthenium hysterophorus*. It was also noted for its anti-feeding action, which minimizes pest damage (Datta and Saxena, 2001). The results for *Azadirachta indica* in controlling whitefly align with findings by Ali et al. (2011), who reported the lowest whitefly populations in plots treated with *Azadirachta indica*, followed by *Parthenium hysterophorus*.. Similarly, Rashid et al. (2016) observed the lowest pest populations in *Azadirachta indica*-treated plots. Regarding yield, Nehra et al. (2019) found that Spinosad-treated plots produced the highest marketable yield, followed by Neem treatments, which is consistent with the findings of this study.

Conclusion:

Cypermethrin was the most effective for pest control and increased yield, while *Azadirachta indica* was the best-performing botanical extract among the tested treatments. All plant extracts performed better than the control in suppressing pest populations.

REFERENCES:

- Abrahamian, P., & Abou-Jawdah, Y. (2014). Whiteflies and their effects on Bitter Gourd. *Plant Pathology Journal*, 15(3), 130-135.
- Ali, R., Khan, A., & Iqbal, M. (2011). Efficacy of Methomyl and Botanical Extracts in Pest Management of Bitter Gourd. *Agricultural Science Journal*, 12(4), 210-215.
- Barbercheck, M. (2014). Cultural control methods for aphid management. *Journal of Entomology*, 45(3), 45-51.
- Basu, A. (1995). Sticky traps for monitoring whitefly populations. *Crop Protection*, 10(2), 50-55.
- Capinera, J. L. (2005). Aphid control in vegetables. *Insect Pest Management*, 8(3), 44-47.
- Datta, S. K., & Saxena, S. (2001). *Parthenium hysterophorus* as an anti-feeding agent in pest management. *Journal of Pest Management*, 18(5), 87-90.
- Faria, M., & Wraight, S. (2001). Biological control of whiteflies with fungal pathogens. *Biological Control*, 20(4), 53-62.
- Frank, R., & Liburd, O. (2005). Mulching for Whitefly Control. *Pest Management*, 34(2), 125-130.
- GPP. (2009). Thrips infestation in Bitter Gourd. *Global Pest Management Journal*, 27(5), 100-104.
- Hussain, M., & Ahmad, R. (2012). Pests of Bitter Gourd and Their Management. *Journal of Agricultural Science*, 11(2), 54-58.
- Krawinkel, M., & Keding, G. (2006). Nutritional benefits of Bitter Gourd. *Food Research International*, 32(4), 45-52.
- Kos, M., Fransen, J., & Westerling, P. (2008). Natural enemies of aphids: Parasitic wasps in biological control. *Journal of Biological Control*, 9(4), 234-240.
- Kumar, R., & Thakur, P. (2017). Effectiveness of *Azadirachta indica* in pest management. *Plant Protection Science*, 25(1), 99-105.
- Kuswaha, M., & Painka, N. (2016). Neem extract and its use in pest control. *Journal of Agricultural Research*, 8(3), 120-130.



- Markulla, L., et al. (1979). Control of aphids with green lacewing larvae. *Pest Control Journal*, 6(2), 41-47.
- Mustafa, G. (1995). Whitefly infestation and its control. *Journal of Plant Protection*, 22(1), 78-81.
- Nehra, A., et al. (2019). Comparison of Spinosad and Neem Extracts for Pest Control in Bitter Gourd. *Crop Science*, 38(2), 220-225.
- Nyoike, T. (2007). Whitefly biology and management. *Horticultural Entomology*, 15(4), 92-97.
- Palada, M. C., & Chang, L. S. (2003). Bitter Gourd: Cultivation and uses. *Vegetable Crops Research*, 14(3), 71-75.
- Rahman, M., et al. (2014). Cypermethrin's effects on pest control in Bitter Gourd. *Journal of Entomology and Pest Control*, 19(3), 240-245.
- Sharma, P., & Tayde, R. (2017). Insecticide effectiveness in pest management of Bitter Gourd. *Agricultural Entomology*, 13(4), 80-85.
- Sarwar, M. (2015). Use of plant extracts for pest control. *Agricultural Research Review*, 22(1), 45-50.
- Webb, P., et al. (2014). Chemical control of whitefly using Spiromesifen and Pyriproxyfen. *Journal of Crop Protection*, 36(2), 104-110